

L# ANSWER 14 OF 419 CAPLUS COPYRIGHT 2002 ACS

AN 2001:927332 CAPLUS

DN 136:53677

TI Preparation of discoloration-free pyrrolidone compositions from amines and γ -butyrolactone

IN Seto, Yoko; Yokoyama, Yoshiyuki; Onoe, Hiroyuki; Takahashi, Kazushige

PA Mitsubishi Chemical Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKXXAF

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PI	JP 2001354646	A2	20011225	JP 2000-177703	20000614
----	---------------	----	----------	----------------	----------

AB Title compns. are prepd. by treatment of γ -butyrolactone (I) contg.

(A) no impurities showing higher UV absorbance than that of I, (B)

.ltoreq.20 impurity compds. detected at .gtoreq.10 ppm by gas chromatog. using a fused silica column chem. bonded with polyethylene glycol, and/or

(C) .ltoreq.15 impurity compds. detected at .gtoreq.10 ppm by gas chromatog. using a fused silica column chem. bonded with

bis(cyanopropyl)methylsilicone, with ammonia, primary, secondary, and/or tertiary amines. Thus, I satisfying the above conditions was autoclaved

with aq. MeNH₂ at 240.degree. for 5 h to give N-methylpyrrolidone contg. .ltoreq.5 ppm ionic impurities (e.g. HCO₂H, AcOH, propionic acid, enanthic acid).

IC ICM C07D201-08

ICS C07D207-267

DT Patent

LA Japanese